
# **HPLC Application**

ID No.: 19704

## LC/MS/MS Analysis of Propoxyphene from urine using Strata-X-Drug B on a Kinetex C18 2.6µm, 50x2.1mm



Retention Time: 2.45 min

#### 2 Propoxyphene

Retention Time: 2.45 min

©2025 Phenomenex Inc. All rights reserved.

Phenomenex products are available worldwide.

www.phenomenex.com.cn

For more information contact your Phenomenex Representative at info@phenomenex.com

ohenomenex

for HPLC Application ID No.: 19704



## LC/MS/MS Analysis of Propoxyphene from urine using Strata-X-Drug B on a Kinetex C18 2.6µm, 50x2.1mm

#### **PRODUCT DESCRIPTION:**

Strata<sup>™</sup>-X-Drug B 33 µm Polymeric Strong Cation, 60 mg / 6 mL, Tubes , 30/Pk

Order No.: 8B-S128-UCH

## SOLID PHASE EXTRACTION (SPE) PRODCEDURE:

**Note:** The solvent volumes shown below are for a 60 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

#### **Condition:**

Load:

Wash:

#### Dry:

10 minutes @ full vacuum

Elute:

#### Final Prep and Analysis:

Evaporate to dryness under a stream of nitrogen at 50 C . Reconstitute in 1 mL of 40% acetonitrile. Inject: 0  $\mu$ L on HPLC Mass Spectrometer (MS) @ amu (ambient)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	рКа	% Rec	%RSC (n=0)
1 Propoxyphene-d11	300			100	
2 Propoxyphene	300			99.21	

**Note:** This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

Phenomenex products are available worldwide.

www.phenomenex.com.cn

For more information contact your Phenomenex Representative at info@phenomenex.com