
HPLC Application ID No.: 19762

Testosterone from Female Human Plasma by LC/MS/MS using Strata-X-A and Kinetex 1.7 µm C18

resusterone	rion rena		Silla Dy LC/	m3/m3 using s	Strata-X-A and Kinetex 1.7 µm C16			
Column:	Kinetex® 1.7	7 µm C18 100 Å,	LC Column 30	x 2.1 mm, Ea				
Dimensions:	30 x 2.1 mm	ID						
Order No:	00A-4475-AN	N			Kinetex			
Elution Type:	Gradient				Ultra-High Performance on Any LC System			
Eluent A:	0.1% Formic	Acid +1 mM Am	m Formate in	Water				
Eluent B:	0.1% Formic	Acid +1 mM Am	m Formate in	ACN				
Gradient	Step No. Time (min) Pct A Pct B							
Profile:	1	0	90	10				
	2	2.5	10	90				
	- 3	3.5	10	90	Products used in this application:			
	4	3.6	90	10				
Flow Rate:	• 0.4 mL/min	5.0	90	10				
	,				🜔 KINETEX.			
Col. Temp.:	ambient		<i>.</i>		TN IN			
Detection:	Mass Spectrometer (MS) @ amu (ambient)							
Detector Info:	Control to the second secon							
19762	hrat—"https://sol	av com/broducte/mace	-chactromatore /lit	m campaign= //114%208	application%20search&utm_source=phenomenex&utm_medium=referral">SCIEX<,			
2.0e4 -		1	1					
1.9e4 -								
1.8e4 -								
1.7e4 -								
1.6e4 - 1.5e4 -								
1.3e4 -								
1.3e4 -								
1.2e4 -								
1.1e4 -								
1.0e4 -								
9000.0 - 8000.0 -								
8000.0 - 7000.0 -								
6000.0								
5000.0 -								
4000.0								



ANALYTES:

- 1 Testosterone Retention Time: 2.62 min
- 2 Testosterone-d3 Retention Time: 2.61 min

©2025 Phenomenex Inc. All rights reserved.

Phenomenex products are available worldwide.

www.phenomenex.com.cn

For more information contact your Phenomenex Representative at info@phenomenex.com

phenomenex

aking with tradition

for HPLC Application ID No.: 19762



Testosterone from Female Human Plasma by LC/MS/MS using Strata-X-A and Kinetex 1.7 µm C18

PRODUCT DESCRIPTION:

Strata[™]-X-A 33 µm Polymeric Strong Anion, 30 mg / 3 mL, Tubes , 50/Pk

Order No.: 8B-S123-TBJ

SOLID PHASE EXTRACTION (SPE) PRODCEDURE:

Note: The solvent volumes shown below are for a 30 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:

Load:

Wash:

Dry:

Dry for 5 min under high vacuum

Elute:

Final Prep and Analysis:

Following evaporation of elution solvent @ 50-55 C under gentle nitrogen stream; Add 50 uL 25% hydroxylamine solution and heat at 60-65 C for 5-10 min, then add 200 uL 5%

Inject: 0 µL on HPLC Mass Spectrometer (MS) @ amu (ambient)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	рКа	% Rec	%RSC (n=0)
1 Testosterone	0				
2 Testosterone-d3	0.5				

Note: This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

Phenomenex products are available worldwide.

www.phenomenex.com.cn

For more information contact your Phenomenex Representative at info@phenomenex.com