

JHW018-hydroxypentyl isomers in Urine using Strata-X-Drug B

Column: Kinetex® 2.6 µm C18 100 Å, LC Column 150 x 3 mm, Ea

Dimensions: 150 x 3 mm ID

Order No: 00F-4462-Y0

Elution Type: Gradient

Eluent A: 10mM ammonium formate

Eluent B: Acetonitrile

Gradient Profile:	Step No.	Time (min)	Pct A	Pct B
	1	0	55	45
	2	7	50	50
	3	7.01	5	95
	4	10	5	95

Flow Rate: 0.6 mL/min

Col. Temp.: ambient

Detection: Mass Spectrometer (MS) @ amu (ambient)

Detector Info: <a target="_blank"

Analyst Note: href="https://sciex.com/products/mass-spectrometers?utm_campaign=2019%20application%20search&utm_source=phenomenex&utm_medium=referral">SCIEX<

Polarity: Positive


CAD: 7.00
CUR: 20.00
GS1: 50.00
GS2: 50.00
IS: 5500.00
TEM: 550.00
ihe: ON
EP 10.00



Products used in this application:



App ID 20926



JHW018-hydroxypentyl isomers in Urine using Strata-X-Drug B

ANALYTES:

- 1 JWH-073-Butanoic acid metabolite
- 2 JWH-018-5-Hydroxypentyl metabolite
- 3 JWH-018-4-Hydroxypentyl metabolite



Sample Preparation Details

for HPLC Application ID No.: 20926

JHW018-hydroxypentyl isomers in Urine using Strata-X-Drug B

PRODUCT DESCRIPTION:

Strata™-X-Drug B 33 µm Polymeric Strong Cation, 60 mg / 3 mL, Tubes , 50/Pk

Order No.: 8B-S128-UBJ

SOLID PHASE EXTRACTION (SPE) PROCEDURE:

Note: The solvent volumes shown below are for a 60 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:

Load:

Sample Hydrolysis: Combine 1 mL Human Urine sample (spiked with analytes at 50 ng/mL), 2 mL of 100 mM sodium acetate buffer, pH 5.0, 25 µL

-D-Glucuronidase (Patella Vulgata from Sigma, 100KU).

Vortex 10-15 secs, followed by incubation for 2 hours in a shaker at 55°C to complete hydrolysis of the glucuronides.

Wash:

Dry:

15 mins under 10-15" of Hg

Elute:

Final Prep and Analysis:

Extraction Protocol

(No need for conditioning or equilibration of the cartridge)

Inject: 10 µL on HPLC Mass Spectrometer (MS) @ amu (ambient)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	pKa	% Rec	%RSC (n=0)
1 JWH-073-Butanoic acid metabolite					
2 JWH-018-5-Hydroxypentyl metabolite					
3 JWH-018-4-Hydroxypentyl metabolite					

Note: This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals.
Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at info@phenomenex.com



Phenomenex products are available worldwide.

www.phenomenex.com.cn

info@phenomenex.com